

# STAGNICOLA RECORDS FROM TURKEY WITH THE DESCRIPTION OF TWO NEW SPECIES, *STAGNICOLA TEKECUS* N. SP. AND *S. KAYSERIS* N. SP. (GASTROPODA: LYMNAEIDAE)

PETER GLÖER<sup>1</sup> & M. ZEKI YILDIRIM<sup>2</sup>

<sup>1</sup>Schulstr. 3, D-25491 Hetlingen, Germany

<sup>2</sup>Süleyman Demirel Üniv. Burdur Eğitim Fak., TR-15100 Burdur, Turkey

*Abstract* Malacological investigations in the counties of Antalya and Isparta (Turkey) have provided new information on the distribution and status of *Stagnicola* species in Asia Minor. By comparing the anatomy of the material collected, two previously undescribed species were recognised:

*Keywords* *Stagnicola tekecus* n. sp., *Stagnicola kayseris* n. sp., *Stagnicola*, *Lymnaeidae*, Turkey

## INTRODUCTION

However, very few segregate *Stagnicola* records exist for southern and eastern Europe, or for Asia Minor. In southern Europe it is believed that the genus *Stagnicola* is represented principally by *S. fuscus* (C. Pfeiffer, 1821), and additionally Bank (Fauna Europaea: www.faunaeur.org) lists *S. corvus* (Gmelin, 1791) for Greece. The only *Stagnicola* species hitherto recorded from Turkey is *S. palustris* (O.F. Müller, 1774) recorded in western Anatolia by Bilgin (1967, 1980) and Yıldırım & Atayeter (2003). Assignment to this taxon is based upon the conchological characters of a specimen illustrated in Bilgin (1967, p. 17, fig. 4). Examination of *Stagnicola* material from Turkey has provided new information on the status and distribution of the species present in that country. In this paper we describe two new species, *Stagnicola tekecus* n. sp. and *S. kayseris* n. sp. and make comparisons with other *Stagnicola* species mentioned from the region.

## MATERIAL AND METHODS

The snails were gathered with a sieve from the banks of the waters. The samples were frozen down to -20 °C over night and after that put in ethanol (75%). This procedure turned out to be as successful as the relaxing technique according to Meier-Brook (in 1976), using pentobarbital. The animals were stretched out completely and could

therefore be dissected without any problems. The tissue is a little more tender through the freezing, as compared to processing with pentobarbital, but still well to handle, however.

The dissections and measurements of the genital organs were carried out using a stereo microscope (Stemi SV 6, Carl Zeiss, Germany), the photographs were made with a digital camera (Nikon D70), and the shell height and shell width were measured using a vernier caliper.

## THE SAMPLING SITES

The sampling sites of *Stagnicola tekecus* n. sp. known up to now are in the counties Antalya and Isparta (Fig. 1), *S. kayseris* n. sp. is only known from the region Kayseri.


*Stagnicola tekecus* n. sp.

**Holotype** Shell 26 mm high, 11.5 mm wide. Zoologisches Museum Hamburg, ZMH 4823 (Fig. 2.1).

**Material examined** We dissected 25 adult specimens of *Stagnicola tekecus* n. sp. and measured 50 shells, which had 5.5 whorls.

**Type locality** Konne, Isparta (37° 52.33' N, 30° 50.38' E, Turkey), IX. 2004 leg. M. Z. Yıldırım.

**Paratypes:** Zoologisches Museum Hamburg, ZMH 4824, Hungarian Natural History Museum, HNHM 94804, Süleyman Demirel Üniv. Burdur Eğitim Fak., Burdur (Turkey), collection of the senior author.



**Figure 1** Sampling sites of *Stagnicola tekecus* n. sp. and *S. kayseris* n. sp. **A:** star: *S. kayseris* n. sp., Kayseri, **B:** *S. tekecus* n. sp. 1: Konne (loc. typ.), 2: Döşemalti.

**Habitat** Calcareous source lake; *S. tekecus* lives there in the muddy shallow water of the lake.

**Derivation of name** The name is given after the region of the locus typicus, Teke.

**Diagnosis** The shell is elongated, conical, light brown to dark brown with 5.5 whorls. It is 23–32 mm high and 11–15 mm wide, the aperture is oval, red brown inside, with a height of 12–16 mm and a width of 8–11 mm (fig. 2.1–2.5). The mantle is unicoloured grey, the head is sprinkled in an irregularly yellowish way (fig. 2.6).

**Anatomy** The ratio of praeputium to phallotheca is 0.38 to 1.78 with the average of 0.81, and it therefore is very variable (fig. 2.7, 2.10). The bursa copulatrix (spermatheca) is spherical, and the spermatheca duct is long and thin (fig. 4.5). The prostata has one fold with two, rarely three, subfolds, which are usually well developed, seldom appearing only as swellings at the basis of the fold (fig. 2.8, 2.9).

#### *Stagnicola kayseris* n. sp.

**Holotype** Shell 16.0 mm high, 7.0 mm wide. Zoologisches Museum Hamburg, ZMH 37564.

**Paratypes** Zoologisches Museum Hamburg, ZMH 37565, Süleyman Demirel Üniv. Burdur Eğitim Fak., Burdur (Turkey), and collection of the senior author.

**Material examined** We dissected and measured the shells of 8 adult specimens from Kayseri-Karpuz Sekisi Basın Gelbula village (38° 40.66' N, 35° 21.34' E, leg. XII. 2002 M. Akman) and 7 adult specimen from Kayseri-Pastirmacilar Park Garden (leg. XII. 2001 M. Z. Yildirim).

**Type locality** Kayseri-Pastirmacilar Park Garden (38° 43.79' N, 35° 29.56' E).

**Habitat** small spring and surroundings.


**Derivation of name** The name is given after the region of the type locality, Kayseri.

**Diagnosis** The shell is elongated, conical, slim, light brown with 5.5 whorls. It is 16–18 mm high and 8–9 mm wide, the aperture is oval, with a height of 0.70–0.75 mm and a width of 4.5 mm. The columellar fold is strong (Fig. 3). The head and the mantle are unicoloured grey.


**Anatomy** The proportion between the praeputium and the phallotheca is ca. 1.0 (n = 15), the bursa copulatrix is spherical, and the spermatheca duct is long and thin, a little widened at the distal end. The prostata has one fold (Fig. 3).

## DISCUSSION

Direct comparison between *Stagnicola tekecus* n. sp., *S. kayseris* n. sp. and the previously mentioned *Stagnicola* species from S-Europe (*S. palus-*



**Figure 2** Shells and anatomy of *Stagnicola tekecus* n. sp. 1-5: Konne (Isparta, Turkey); 6: Mantle; 7: Praeputium and phallosome; 8-9: Prostata; 10: Genital organs. ag = albumin gland; bc = bursa copulatrix; bd = bursa duct; cp = corpus pyriforme; hd = hermaphroditic duct; m = muscle; ng = nidamental gland; pht = phallosome; prp = praeputium; u = uterus; v = vagina; vd = vas deferens.



**Figure 3** *Stagnicola kayseris* n. sp. – bc = bursa copulatrix, bd = bursa duct, pht = phallosome, prp = praeputium, pvd = provaginal duct, pr = prostata, prp = praeputium, vd = vas deferens.



**Figure 4** *Stagnicola* species: 1, 1a *S. fuscus* (Mórfion, Thesprotia, Greece), 2, 2a *S. corvus* (Treibtal, N-Germany), 3, 3a *S. palustris* (Peenetal, N-Germany), 4 *S. kayseris* (Kayseri, Turkey), 5 *S. tekecus* (Konne, Turkey). Magnification: shells 2x, all others 10x.

*tris*, *S. fuscus*, *S. corvus*) reveals the difference of the morphology of the shells as well as the male and female sex tract (see fig. 4). While the shells of the species living in Central Europe have a large phenotypical plasticity (Jackiewicz 1998, Glöer 2002), the shells of the newly described species do not.

**DIFFERENTIAL DIAGNOSIS**

While *Stagnicola corvus* differs from all other known species in the number of prostata folds, the anatomies of *S. palustris*/*S. kayseris* respectively *S. fuscus*/*S. tekecus* in some respect look similar. But *S. fuscus* has a short phallotheca

and so this feature is in significant contrast to *S. tekecus* n. sp.

*S. kayseris* has a broad provaginal duct, which is shorter than in *S. palustris*, and the spermathecal duct is a little widened at the distal end. The shells of *S. kayseris* differ significantly from the other species mentioned, e. g. the strong columellar fold.

**ACKNOWLEDGEMENTS**

We would like to express our gratitude to the two unknown referees for helpful comments, and Dr David Walker (Hamburg) for helping to polish our english.

## REFERENCES

- BILGIN F 1967 İzmir Civari Tath Sularında Yaşayan Gastropodlar Üzerinde Sistemik ve Ekolojik Araştırmalar (Ecological and Systematical research on the living gastropods in fresh waters in the vicinity of İzmir. *Scientific reports of the Faculty of Science Ege University* 36: 1-54, 2 pl. Bornova.
- BILGIN F 1980 Batı Anadolu'nun bazı önemli tath sularından toplanan Mollusca türlerini sistematiği ve dağılışı (Systematics and distribution of Mollusca species collected from some fresh waters of West Anatolia). *Diyarbakir Üniversitesi Tıp Fakültesi Dergisi* 8 (2) (Supplement): 64 pp. Diyarbakir.
- GLÖER P 2002 Die Süßwassergastropoden Nord- und Mitteleuropas. *Die Tierwelt Deutschlands* 73: 327 pp. (ConchBooks) Hackenheim.
- JACKIEWICZ M 1998 European species of the family Lymnaeidae (Gastropoda: Pulmonata: Basommatophora). *Genus - International journal of invertebrate taxonomy* 9 (1): 1-93. Poznan.
- MEIER-BROOK C 1976 An improved relaxing technique for molluscs using pentobarbital. *Malacological Review* 9: 115-117. Whitmore Lake.
- YILDIRIM MZ & ATAYETER Y 2003 General geographic and malakologic features of lakes in the Lakes Region (Turkey). *International Symposium of fisheries and Zoology*. 23-26 October, Istanbul.